


# Nevolat® (liraglutide)

## Dosing Guide

Your Nevolat® treatment will start at a low dose (0.6 mg), which will gradually increase over the first five weeks of use. Starting from Week 5 of treatment, you will be taking Liraglutide at a dose of 3.0 mg every day. This is the highest dose of Nevolat® that you will take during your treatment – do not increase the dose any further.



### Starting Dose

#### Nevolat® Treatment Guide: First 8 Weeks

##### Daily Dosage:

Refer to the colour of each square to determine the dosage required for that day. each dose is maintained for at least one week, but may be longer before increasing if advised by the HCP

**Pen Changes:** Squares outlined in pink indicate when you need to change Nevolat® pens.

##### Tracking:

Check off each day in the bottom right-hand corner of the square to make sure you stay up to date with your daily dosing.

### Full Dose

1 0.6mg	2 0.6mg	3 0.6mg	4 0.6mg	5 0.6mg	6 0.6mg	7 0.6mg
8 1.2mg	9 1.2mg	10 1.2mg	11 1.2mg	12 1.2mg	13 1.2mg	14 1.2mg
15 1.8mg	16 1.8mg	17 1.8mg	18 1.8mg	19 1.8mg	20 1.8mg	21 1.8mg
22 2.4mg	23 2.4mg	24 2.4mg	25 2.4mg	26 2.4mg	27 2.4mg	28 2.4mg
29 3.0mg	30 3.0mg	31 3.0mg	32 3.0mg	33 3.0mg	34 3.0mg	35 3.0mg
36 3.0mg	37 3.0mg	38 3.0mg	39 3.0mg	40 3.0mg	41 3.0mg	42 3.0mg
43 3.0mg	44 3.0mg	45 3.0mg	46 3.0mg	47 3.0mg	48 3.0mg	49 3.0mg
50 3.0mg	51 3.0mg	52 3.0mg	53 3.0mg	54 3.0mg	55 3.0mg	56 3.0mg



Always use this medicine exactly as your doctor has told you. Check with your doctor, pharmacist or nurse if you have any questions about taking Nevolat®.

Date of preparation: March 2025.  
Reference: 000741646.

Adverse events should be reported. Reporting forms and information can be found at [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard). Adverse events should also be reported to Zentiva via email to [UKmedinfo@zentiva.com](mailto:UKmedinfo@zentiva.com) or via phone on 0800 090 2408.