


Zegluxen® (liraglutide)

Dosing Guide

To help minimise any digestive issues, Zegluxen® is started at a lower dose and gradually increased over time. The starting dose is **0.6 mg once a day**, for at least one week. Your doctor will tell you when to increase it to 1.2 mg once a day. Your doctor may tell you to further increase the dose to 1.8 mg once a day, if your blood sugar level is not adequately controlled with a dose of 1.2 mg.



Zegluxen® Treatment Guide: First 8 Weeks

Daily Dosage:

Refer to the color of each square to determine the dosage required for that day.

Add your personalised dose instructions as advised by your doctor.

Tracking:

Check off each day in the bottom right-hand corner of the square to make sure you stay up to date with your daily dosing.

1 0.6mg	2 0.6mg	3 0.6mg	4 0.6mg	5 0.6mg	6 0.6mg	7 0.6mg
8 1.2mg	9 1.2mg	10 1.2mg	11 1.2mg	12 1.2mg	13 1.2mg	14 1.2mg
15	16	17	18	19	20	21
22	23	24	25	26	27	28
29	30	31	32	33	34	35
36	37	38	39	40	41	42
43	44	45	46	47	48	49
50	51	52	53	54	55	56



Always use this medicine exactly as your doctor has told you. Check with your doctor, pharmacist or nurse if you have any questions about taking Zegluxen®.
Date of preparation: March 2025.
Reference: 000741647

Adverse events should be reported. Reporting forms and information can be found at www.mhra.gov.uk/yellowcard. Adverse events should also be reported to Zentiva via email to UKmedinfo@zentiva.com or via phone on **0800 090 2408**.